

Fixed Point Convergence

Math 426

University of Alaska Fairbanks

September 16, 2020

Secant Method: Rate of Convergence

$$f(x) = x^2 - 2, \quad x_1 = 1, \quad x_2 = 1.1$$

$$e_3 \approx 4 \times 10^{-1}$$

$$e_4 \approx 3 \times 10^{-1}$$

$$e_5 \approx 6 \times 10^{-2}$$

$$e_6 \approx 8 \times 10^{-3}$$

$$e_7 \approx 2 \times 10^{-4}$$

$$e_8 \approx 4 \times 10^{-7}$$

$$e_9 \approx 2 \times 10^{-11}$$

$$e_{10} \approx 4 \times 10^{-16}$$

$$\frac{|e_{k+1}|}{|e_k|^\alpha} \rightarrow C > 0$$

$$|e_{k+1}| \approx C |e_k|^\alpha$$

bisection $\alpha = 1$

Newton: $\alpha = 2$

Secant Method: Rate of Convergence


Text:

Secant

$$\lim_{k \rightarrow \infty} \frac{|e_{k+1}|}{|e_k e_{k-1}|} = C$$

Newton:

$$\lim_{k \rightarrow \infty} \frac{|e_{k+1}|}{|e_k|^2} = C$$



Secant Method: Rate of Convergence

Text:

$$\lim_{k \rightarrow \infty} \frac{|e_{k+1}|}{|e_k e_{k-1}|} = C$$

Newton:

$$\lim_{k \rightarrow \infty} \frac{|e_{k+1}|}{|e_k|^2} = C$$

Suppose $e_{k+1} \sim a e_k^\alpha$.

$$\frac{|e_{k+1}|}{|e_k|^\alpha} \rightarrow a$$

Secant Method: Rate of Convergence

Text:

$$\lim_{k \rightarrow \infty} \frac{|e_{k+1}|}{|e_k e_{k-1}|} = C$$

Newton:

$$\lim_{k \rightarrow \infty} \frac{|e_{k+1}|}{|e_k|^2} = C$$


Suppose $e_{k+1} \sim a e_k^\alpha$.

$e_k \sim a e_{k-1}^\alpha \Rightarrow$

$a^{-1} e_k \sim e_{k-1}^\alpha$
 $a^{-1/\alpha} e_k^{1/\alpha} \sim e_{k-1}$

$$\frac{|e_{k+1}|}{|e_k e_{k-1}|} \sim \frac{a |e_k|^\alpha}{|e_k| |e_{k-1}|} \sim \frac{a |e_k|^\alpha}{a^{-1/\alpha} |e_k| |e_k|^{1/\alpha}} \sim c |e_k|^{\alpha-1-1/\alpha}$$

$|e_k| |e_{k-1}|$



$e_k \rightarrow 0$

Secant Method: Rate of Convergence

Text:

$$\lim_{k \rightarrow \infty} \frac{|e_{k+1}|}{|e_k e_{k-1}|} = C$$

Newton:

$$\lim_{k \rightarrow \infty} \frac{|e_{k+1}|}{|e_k|^2} = C$$

Suppose $e_{k+1} \sim a e_k^\alpha$.

$$\frac{|e_{k+1}|}{|e_k e_{k-1}|} \sim \frac{a |e_k|^\alpha}{|e_k| |e_{k-1}|} \sim \frac{a |e_k|^\alpha}{a^{-1/\alpha} |e_k| |e_k|^{1/\alpha}} \sim c |e_k|^{\alpha-1-1/\alpha}.$$

$$e_k \rightarrow 0 \implies \alpha^2 - \alpha - 1 = 0$$

$$\alpha - 1 - \frac{1}{\alpha} = 0$$

$$\alpha^2 - \alpha - 1 = 0$$

Secant Method: Rate of Convergence

Text:

$$\lim_{k \rightarrow \infty} \frac{|e_{k+1}|}{|e_k e_{k-1}|} = C$$

Taylor's Theorem

Newton:

$$\lim_{k \rightarrow \infty} \frac{|e_{k+1}|}{|e_k|^2} = C$$

$a < 0$

$$e_{k+1} \sim a e_k^2$$

Suppose $|e_{k+1}| \sim a |e_k|^\alpha$

$$\frac{|e_{k+1}|}{|e_k e_{k-1}|} \sim \frac{a |e_k|^\alpha}{|e_k| |e_{k-1}|} \sim \frac{a |e_k|^\alpha}{a^{-1/\alpha} |e_k| |e_k|^{1/\alpha}} \sim c |e_k|^{\alpha-1-1/\alpha}$$

$$\frac{|e_{k+1}|}{|e_k|^\alpha} \rightarrow C$$

$$e_k \rightarrow 0 \implies \alpha^2 - \alpha - 1 = 0$$

$$\alpha = \frac{1 \pm \sqrt{1+4}}{2} = \frac{1 \pm \sqrt{5}}{2}$$

$$\frac{1 - \sqrt{5}}{2} < 0$$

$$e_k \rightarrow 0 \implies \alpha = \frac{1 + \sqrt{5}}{2} \approx 1.6$$

Convergence of Newton's Method

We showed that **if** Newton's method converges and $f'(x_*) \neq 0$ then

$$\frac{|e_{k+1}|}{|e_k|^2} \rightarrow \frac{f''(x_*)}{2f'(x_*)} \cdot C$$

But we never showed that it would actually converge.

Convergence of Newton's Method

We showed that **if** Newton's method converges and $f'(x_*) \neq 0$ then

$$\frac{|e_{k+1}|}{|e_k|^2} \rightarrow \frac{f''(x_*)}{2f'(x_*)}.$$

But we never showed that it would actually converge.

Strategy: formulate Newton's method as a fixed point method, determine conditions under which fixed point methods converge, and apply to Newton's method.

Fixed point methods

Want to solve

$$\Phi(x) = x.$$

$$\underline{\Phi}(x_k) = x_{k+1}$$

Such a point is called a **fixed point** of Φ .

For Newton's method,

$$\Phi(x) = x - \frac{f(x)}{f'(x)}$$

$$\underline{\Phi}(x) = x$$

Fixed point methods

Want to solve

$$\Phi(x) = x.$$

Such a point is called a **fixed point** of Φ .

For Newton's method,

$$\Phi(x) = x - \frac{f(x)}{f'(x)}$$

If $\Phi(x_*) = x_*$,

$$x_* = x_* - \frac{f(x_*)}{f'(x_*)}$$

so

$$f(x_*) = 0.$$

$$0 = - \frac{f(x_*)}{f'(x_*)}$$

Fixed point methods

Want to solve

$$\Phi(x) = x.$$

Such a point is called a **fixed point** of Φ .

For Newton's method,

$$|f(x_k)| < f_{tol}$$

$$\Phi(x) = x - \frac{f(x)}{f'(x)}$$

If $\Phi(x_*) = x_*$,

$$x_* = x_* - \frac{f(x_*)}{f'(x_*)}$$

so

$$f(x_*) = 0.$$

If $f(x_*) = 0$ then $\Phi(x_*) = x_*$ (assuming $f'(x_*) \neq 0$.)

N_{max}

• f_{tol}

• x_{tol}

$$|x_k - x_{k-1}|$$

$$< x_{tol}$$

$$\Phi(x_*) = x_* - \frac{f(x_*)}{f'(x_*)}$$

$$= x_* - \frac{0}{f'(x_*)}$$

$$= x_*$$

Fixed point methods

Want to solve

$$\Phi(x) = x.$$

Such a point is called a **fixed point** of Φ .

For Newton's method,

$$\Phi(x) = x - \frac{f(x)}{f'(x)}$$

If $\Phi(x_*) = x_*$,

$$x_* = x_* - \frac{f(x_*)}{f'(x_*)}$$

so

$$f(x_*) = 0.$$

If $f(x_*) = 0$ then $\Phi(x_*) = x_*$ (assuming $f'(x_*) \neq 0$).

The fixed points of Φ are exactly the roots of f (up to legalese).

Visualizing Fixed Point Iteration

Suppose $\Phi(x) = x^2$. What are the fixed points?

$$x = 1$$


$$x = 0$$

$$\Phi(1) = 1^2 = 1$$

$$\Phi(0) = 0^2 = 0$$


Visualizing Fixed Point Iteration

Suppose $\Phi(x) = x^2$. What are the fixed points?



Visualizing Fixed Point Iteration

Suppose $\Phi(x) = x^2$. What are the fixed points?



Visualizing Fixed Point Iteration

Suppose $\Phi(x) = x^2$. What are the fixed points?


$$x = 0 \quad x = 1$$

$$(x_k, \Phi(x_k)) \rightarrow (\Phi(x_k), \Phi(x_k)) = (x_{k+1}, x_{k+1}) \rightarrow (x_{k+1}, \Phi(x_{k+1}))$$


x_{k+2}

Two cases for $\Phi(x) = x^2$


$$\phi(x) = x$$



Two cases for $\Phi(x) = x^2$



$$\Phi(x) = x/2$$



$$x_2 = \Phi(x_1) = \frac{1}{2} x_1$$

$$x_3 = \Phi(x_2) = \frac{1}{2} x_2 = \frac{1}{4} x_1$$

$$\Phi(x) = 3x$$

$$\Phi(x) = cx$$

$$x_k = c^{k-1} x_1$$

$|c| < 1 \rightarrow$ converge
 $|c| > 1 \rightarrow$ diverge


x_1

$$x_1 = 5$$

$$x_2 = 3 \cdot 5$$

$$x_3 = 3^2 \cdot 5$$

$$x_k = 3^{k-1} \cdot 5$$




$$\Phi(x) = -x/2$$

$$x_2 = -\frac{1}{2}x_1$$


$$x_3 = +\frac{1}{4}x_1$$

$$x_4 = -\frac{1}{8}x_1$$



$$\Phi(x) = -2x$$

$$|-2| = 2 > 1$$



$$\Phi(x) = -2x$$

