

4.3? Partial Derivatives

$$f(x, y) = e^x \cos(x^3 y^2)$$

$$\frac{\partial f}{\partial x} = e^x \cos(x^3 y^2) - e^x \sin(x^3 y^2) \cdot (3x^2 y^2)$$

↑

partial derivative

of f with respect to x


$$\frac{d}{dy} e^7 \cos(7^3 y^2) = -e^7 \sin(7^3 y^2) \cdot (2 \cdot 7^3 y)$$

$$\frac{\partial}{\partial y} e^x \cos(x^3 y^2) = -e^x \sin(x^3 y^2) \cdot (2x^3 y)$$

$$P = \frac{nRT}{V}$$

$$l = nR \frac{T}{V}$$

$$V = (nR) T$$





$$R = 0.082 \frac{\text{atm l}}{\text{mol K}}$$

$$P = 0.82 \text{ atm}$$

$$P = \frac{0.042 \cdot 300}{V}$$

$$\frac{dP}{dV} = - \frac{0.042 \cdot 300}{V^2} \quad \checkmark \quad \frac{\text{atm}}{\text{l}}$$

$$\left. \frac{dP}{dV} \right|_{V=30\text{l}} = -0.027 \frac{\text{atm}}{\text{l}}$$



$$P = \frac{0.092 \cdot 300}{V}$$

$$\left. \frac{dP}{dV} \right|_{V=30L} = -0.027 \frac{\text{atm}}{L}$$

$$\frac{\partial p}{\partial V} = -\frac{0.082 \cdot T}{V^2}$$

$$\left(p = \frac{0.082 T}{V} \right)$$



If the temp of the gas is T and the volume is V and we change the volume the pressure changes at a rate of

$$-\frac{0.082 \cdot T}{V^2}$$

$$T=300, V=20 \Rightarrow \frac{\partial p}{\partial V} = -0.0615 \text{ atm/l}$$

$$T=300, V=30 \Rightarrow \frac{\partial p}{\partial V} = -0.027 \text{ atm/l}$$

Ballpark: how much should I increase the volume from 30L to decrease the pressure by 0.1 atm.

(0.92 atm at $T=300\text{K}$, $V=30\text{L}$)

$$-0.027 \frac{\text{atm}}{\text{L}} \cdot \Delta V \text{L} = -0.1 \text{ atm}$$

$$\Delta V = \frac{-0.1}{-0.027} = 3.7 \text{ L}$$

$$P = \frac{0.092 \cdot 300}{33.7} = 0.73 \text{ atm}$$

Summary: Given $f(x, y)$

$\frac{\partial f}{\partial x}(x_0, y_0)$ is the rate of change of f with respect to x at $x = x_0, y = y_0$.

Second partial derivatives

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \frac{\partial}{\partial x} f = f_{xx}$$

$$\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial y} \frac{\partial}{\partial x} f = f_{xy}$$

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial x} \frac{\partial}{\partial y} f = f_{yx}$$

Pure magic:

$$f(x, y) = x^3 y^2 - x \ln(y)$$

$$f_x = 3x^2 y^2 - \ln(y)$$

$$f_y = 2x^3 y - \frac{x}{y}$$

$$f_{xy} = 6x^2 y - \frac{1}{y}$$

$$f_{yx} = 6x^2 y - \frac{1}{y}$$

Up to legalese $f_{xy} = f_{yx}$ always.

$$f_{xy} = f_{yx} = f_{yx}$$