



$$\vec{F} = \langle x^2, x^4 \rangle$$



$P$	$\vec{F}$
$(0,0)$	$\langle 0,0 \rangle$
$(1,1)$	$\langle 1,1 \rangle$
$(1,2)$	$\langle 1,2 \rangle$
$(2,1)$	$\langle 4,2 \rangle$

$$\langle x, y \rangle$$



$$\langle x, y \rangle$$


$$\langle x, y \rangle$$

$$- \left\langle \frac{x}{\sqrt{x^2+y^2}}, \frac{y}{\sqrt{x^2+y^2}} \right\rangle$$



FTC

$$\int_a^b f'(t) dt = f(b) - f(a)$$



$$\vec{F} = \vec{\nabla} f$$

Fund. Thm of Line Integrals

$$\int_C \vec{\nabla} f \cdot d\vec{r} = f(\vec{r}(b)) - f(\vec{r}(a))$$

$$\vec{\nabla} T \cdot \vec{r}'$$

$$\vec{r}(t) = \langle x(t), y(t) \rangle \quad a \leq t \leq b$$

$$\vec{r}'(t) = \langle x'(t), y'(t) \rangle$$


$$\int_C \vec{\nabla} f \cdot d\vec{r} = \int_a^b \nabla f \cdot \vec{r}'(t) dt$$

$$= \int_a^b \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right\rangle \cdot \langle x', y' \rangle dt$$

$$= \int_a^b \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} dt$$

$$= \int_a^b \frac{d}{dt} f(\vec{r}(t)) dt$$


$$= f(\vec{r}(b)) - f(\vec{r}(a))$$



$$\vec{F} = \vec{\nabla} f \quad \text{potential for } \vec{F}$$

$\vec{F}$   $\rightarrow$  conservative vector field


$\int_C \vec{\nabla} f \cdot d\vec{r}$  depends only on the endpoints (with orientation)



$$\int_C \vec{\nabla} f \cdot d\vec{r} = f(q) - f(p)$$

$$\int_{\hat{C}} \vec{\nabla} f \cdot d\vec{r} = f(p) - f(q)$$

$$\int_C \vec{\nabla} f \cdot d\vec{r} = f(q) - f(p)$$



$$\int_C \vec{\nabla} f \cdot d\vec{r} = f(p) - f(p) = 0$$

Three properties of conservative vector fields  $\vec{F} = \langle P, Q \rangle$

a) the integral over a curve depends only on the end points


b) the integral around a loop is zero

$$c) \frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}, \quad \frac{\partial P}{\partial z} = \frac{\partial R}{\partial x}, \quad \frac{\partial Q}{\partial z} = \frac{\partial R}{\partial y} \quad \langle P, Q, R \rangle$$

$$\vec{F} = \langle P, Q \rangle = \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right\rangle$$

$$\frac{\partial P}{\partial y} = \frac{\partial}{\partial y} \frac{\partial f}{\partial x} = \frac{\partial}{\partial x} \frac{\partial f}{\partial y} = \frac{\partial}{\partial x} Q = \frac{\partial Q}{\partial x}$$


---



$$\vec{F} = -\frac{MmG}{|\vec{x}|^2} \frac{\vec{x}}{|\vec{x}|} = -\frac{MmG}{|\vec{x}|^3} \vec{x}$$

Claim:  $\vec{F} = \nabla f$

$$f = \frac{MmG}{|\vec{x}|}$$



$$\text{work} \int_C \vec{F} \cdot d\vec{r} = f(q) - f(p) \\ = M_m G \left( \frac{1}{|\vec{x}(q)|} - \frac{1}{|\vec{x}(p)|} \right)$$


$$|\vec{x}(p)| = R_e \text{ radius of earth} \\ = 6.365 \times 10^6 \text{ m}$$

$$|\vec{x}(q)| = R_o \text{ radius in orbit} \\ = 6.765 \times 10^6 \text{ m}$$

$$E = -296 \text{ MJ}$$


$$M G = 3.98 \times 10^{14}$$

$$m = 80 \text{ kg}$$



$\int_C \vec{F} \cdot d\vec{r}$  depends only on endpoints

$\vec{F}$  Suppose  $\oint_C \vec{F} \cdot d\vec{r} = 0$  where  $C$  is a loop



Want to show

$\int_C \vec{F} \cdot d\vec{r}$  depends only on endpoints





$$\int_C \vec{F} \cdot d\vec{r} \approx \int_{C_1} \vec{F} \cdot d\vec{r}$$


$$\int_C \vec{F} \cdot d\vec{r} - \int_{C_1} \vec{F} \cdot d\vec{r} \approx 0$$

$$\int_{C_2} \vec{F} \cdot d\vec{r} = 0$$

$$\int_C \vec{F} \cdot d\vec{r} = \int_{C_1} \vec{F} \cdot d\vec{r} + \int_{C_2} \vec{F} \cdot d\vec{r} = 0$$



$$\int_C \vec{F} \cdot d\vec{r} = \int_{C_1} \vec{F} \cdot d\vec{r} + \int_{C_2} \vec{F} \cdot d\vec{r}$$



$$\int_C \vec{F} \cdot d\vec{r} = \int_{C_1} \vec{F} \cdot d\vec{r} - \int_{C_2} \vec{F} \cdot d\vec{r}$$

$$= 0.$$


---

$$\vec{F}(x, y) = \left\langle \underbrace{3 + 2xy}_P, \underbrace{x^2 - 3y^2}_Q \right\rangle$$

$$\frac{\partial P}{\partial y} = 2x \quad \frac{\partial Q}{\partial x} = 2x$$

same.

$$\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$$

Is  $\vec{F}$  conservative?

f

$$\frac{\partial f}{\partial x} = 3 + 2xy$$

$$\frac{\partial f}{\partial y} = x^2 - 3y^2$$

$$\frac{\partial f}{\partial x} = 3 + 2xy$$

$$f = 3x + x^2y + h(y)$$

$$\frac{\partial f}{\partial y} = 0 + x^2 + h'(y)$$

$$h'(y) = -3y^2 \quad h(y) = -y^3$$

$$f(x, y) = 3x + x^2y - y^3$$