

The Weierstrass Approximation Theorem states that a function in  $C[a, b]$  can be uniformly approximated by a polynomial. One way of expressing this fact is that given  $f \in C[a, b]$  and  $\epsilon > 0$ , there exists  $p \in P[a, b]$  such that  $|f(x) - p(x)| \leq \epsilon$  for every  $x \in [a, b]$ . Using the vocabulary of norms, this is equivalent to

$$\|f - p\|_\infty \leq \epsilon.$$

The same idea can also be expressed in terms of the closure of  $P[a, b]$  in  $C[a, b]$ . Recall that given a set  $A$  in a metric space  $M$ ,  $x \in \bar{A}$  if and only if for every  $\epsilon > 0$ ,  $B_\epsilon(x) \cap A \neq \emptyset$ . Hence the Weierstrass Approximation Theorem asserts that  $C[a, b] \subseteq \overline{P[a, b]}$ . But of course  $\overline{P[a, b]} \subseteq C[a, b]$ . Hence we have arrived at a concise statement of the theorem.

**Theorem 1: (Weierstrass Approximation Theorem)**  $\overline{P[a, b]} = C[a, b]$ , where closure is taken with respect to the uniform norm.

You are already familiar with the idea of writing certain functions as power series. For example,

$$\sin(x) = \sum_{n=0}^{\infty} \frac{(-1)^{n+1} x^{2n+1}}{n!}.$$

This series converges pointwise on all of  $\mathbb{R}$  (verify this with the ratio test) and therefore uniformly on any fixed interval  $[-R, R]$ . (Recall Theorem 10.10) Hence, given any  $\epsilon > 0$ , we can find an  $N$  such that

$$\left| \sin(x) - \sum_{n=0}^N \frac{(-1)^{n+1} x^{2n+1}}{n!} \right| \leq \epsilon$$

for every  $x \in [-\pi, \pi]$ . So  $\sin$  can be approximated uniformly by polynomials on  $[-\pi, \pi]$ . But functions that can be written as power series are special; in particular they are infinitely differentiable – this is a consequence of Theorem 10.10.


The remarkable part about the Weierstrass Approximation Theorem is that every continuous function, even the non-differentiable ones, can be uniformly approximated by polynomials. Interestingly, the proof of this fact can be reduced to showing that just one non-smooth function, the absolute value function  $\text{abs}$ , can be uniformly approximated by polynomials.

**Proposition 2:**  $\text{abs} \in \overline{P[-1, 1]}$ .

Supposing for the moment we have proved this result, let's see how this results in a fairly easy proof of Theorem 1. First, we show that any translate of the absolute value function is in  $\overline{P[0, 1]}$ . We define

$$\text{abs}_a(x) = |x - a|.$$

**Lemma 3:** For any  $a \in \mathbb{R}$ ,  $\text{abs}_a \in \overline{P[0, 1]}$ .

Approximating  $\text{abs}_a$  on  $[0, 1]$ .

*Proof.* If  $a \leq 0$  or  $a \geq 1$ ,  $\text{abs}_a$  is linear on  $[0, 1]$  and hence in  $P[0, 1]$ .

Suppose  $0 < a < 1$  and let  $\epsilon > 0$ . Let  $p$  be a polynomial such that

$$|p(x) - \text{abs}(x)| < \epsilon$$

for every  $x \in [-1, 1]$ . Define  $q(x) = p(x - a)$ , so  $q$  is a polynomial. Then

$$\begin{aligned} \sup_{x \in [0, 1]} |q(x) - \text{abs}_a(x)| &= \sup_{x \in [0, 1]} |p(x - a) - |x - a|| \\ &= \sup_{x \in [-a, 1-a]} |p(x) - |x|| \\ &\leq \sup_{x \in [-1, 1]} |p(x) - \text{abs}(x)| \leq \epsilon. \end{aligned}$$

Hence  $\|q - \text{abs}_a\|_{C[0, 1]} \leq \epsilon$ . Since  $q$  is a polynomial and  $\epsilon > 0$  is arbitrary,  $\text{abs}_a \in \overline{P[0, 1]}$ .  $\square$

A function  $f \in C[0, 1]$  is called piecewise linear if there is a partition  $0 = x_0 < x_1 < \dots < x_n = 1$  such that the restriction of  $f$  to each interval  $[x_{k-1}, x_k]$  is linear; we denote by  $\text{PL}[0, 1]$  the collection of all such functions. Clearly any linear combination of functions of the form  $\text{abs}_a$  belongs to  $\text{PL}[0, 1]$ . We now show that these functions span all of  $\text{PL}[0, 1]$ .

**Proposition 4:** Let  $f \in \text{PL}[0, 1]$ , and let  $0 = x_0 < x_1 < \dots < x_n = 1$  be a partition such that  $f$  is linear on each interval  $I_k = [x_{k-1}, x_k]$ . Then  $f$  is a linear combination of the functions 1 and  $\{\text{abs}_{x_k} : 0 \leq k \leq n\}$ .


*Proof.* Let  $S = \text{span}\{\text{abs}_{x_k} : 0 \leq k \leq n\}$ . Notice that

$$\text{abs}_{x_0}(x) + \text{abs}_{x_n}(x) = x + (1 - x) = 1.$$

Hence the constants belong to  $S$ .

For  $0 \leq k \leq 1$ , let

$$R_k(x) = \frac{1}{2} (\text{abs}_{x_k}(x) + (x - x_k)).$$

The functions  $R_k$ ,  $J_k$ , and  $H_k$ .

Then  $R_k$  is a linear combination of 1,  $\text{abs}_{x_0}$ , and  $\text{abs}_{x_k}$  and hence  $R_k \in S$ .

Notice that  $R_k(x) = 0$  if  $x \leq x_k$  and  $R_k(x) = x - x_k$  otherwise. For  $1 \leq k \leq n$  let

$$J_k = \frac{R_k - R_{k-1}}{x_k - x_{k-1}},$$

and let  $J_0 = 1$  and  $J_{n+1} = 0$ . Then each  $J_k \in S$  and

$$J_k(x_j) = \begin{cases} 0 & j < k \\ 1 & j \geq k. \end{cases}$$

Finally, let  $H_k = J_k - J_{k+1}$  for  $0 \leq k \leq n$ . Then  $H_k \in S$  for each  $k$ , and

$$H_k(x_j) = \begin{cases} 1 & k = j \\ 0 & k \neq j. \end{cases}$$

Hence

$$\sum_{k=0}^n f(x_k) H_k$$

is a piecewise linear function that agrees with  $f$  at each point  $x_k$ . We conclude that

$$f = \sum_{k=0}^n f(x_k) H_k.$$

Since each  $H_k \in S$ , we conclude that  $f \in S$ . □

We have seen that each  $\text{abs}_a \in \overline{P[0,1]}$  and that each  $f \in PL[0,1]$  is a linear combination of functions  $\text{abs}_a$ . To show that  $PL[0,1] \subseteq \overline{P[0,1]}$  we now take advantage of the idea that the metric space and the vector space structures of a normed vector space are compatible.

**Proposition 5:** Let  $X$  be a normed linear space and let  $W$  be a subspace of  $X$ . Then  $\overline{W}$  is a subspace of  $X$ .

*Proof.* Let  $x, y \in \overline{W}$ . Let  $(x_n)$  and  $(y_n)$  be sequences in  $W$  converging to  $x$  and  $y$ . Then  $\|(x + y) - (x_n + y_n)\| \leq \|x - x_n\| + \|y - y_n\|$  and therefore  $(x_n + y_n) \rightarrow (x + y)$ . Hence  $x + y \in \overline{W}$ . Similarly,  $\alpha x_n \rightarrow \alpha x$  and hence  $\alpha x \in \overline{W}$ . So  $\overline{W}$  is a subspace.  $\square$

We can now prove the Weierstrass Approximation Theorem, at least for the domain  $[0, 1]$ .

**Proposition 6:**  $C[0, 1] = \overline{P[0, 1]}$ .

*Proof.* Proposition 5 implies that  $\overline{P[0, 1]}$  is a subspace of  $C[0, 1]$  since  $P[0, 1]$  is. Suppose  $f \in PL[0, 1]$ . Proposition 4 shows that  $f$  can be written as a finite linear combination of functions  $\text{abs}_a$ , and Proposition 3 implies that each  $\text{abs}_a \in \overline{P[0, 1]}$ . Since  $\overline{P[0, 1]}$  is a subspace, we conclude that  $f \in \overline{P[0, 1]}$  and hence  $PL[0, 1] \subseteq \overline{P[0, 1]}$ . Consequently  $\overline{PL[0, 1]} \subseteq \overline{P[0, 1]}$ . From the proof of Carothers 11.2 it follows that  $C[0, 1] = \overline{PL[0, 1]}$ . Hence  $\overline{P[0, 1]} = C[0, 1]$ .  $\square$

**Exercise 1:** Use Proposition 6 to prove the Weierstrass Approximation Theorem for an arbitrary interval  $[a, b]$ . Hint: Given  $f \in C[a, b]$ , define  $g(x) = f(a + x(b - a))$ . Approximate  $g$  in  $C[0, 1]$  by  $p \in P[0, 1]$ , and define  $q(x) = p((x - a)/(b - a))$ .

It remains to prove Proposition 2, which we do now.

*Proof.* For  $0 \leq x \leq 1$ , define  $P_0(x) = 0$  and for  $k \geq 0$  define

$$P_{k+1}(x) = P_k(x) + \frac{x - P_k^2}{2}.$$

We claim that  $0 \leq P_k(x) \leq \sqrt{x}$  for every  $k \geq 0$  and that  $P_{k+1} \geq P_k$  for every  $k$ . This is certainly true for  $k = 0$ . Suppose  $0 \leq P_k(x) \leq \sqrt{x}$ . Then

$$P_{k+1}(x) = P_k(x) + \frac{x - P_k^2}{2} \geq P_k(x)$$

so  $P_{k+1}(x) \geq 0$ . But also, since  $0 \leq P_k(x) \leq \sqrt{x} \leq 1$ , we have

$$P_{k+1}(x) = P_k(x) + \frac{x - P_k^2}{2} = P_k(x) + \frac{1}{2}(\sqrt{x} + P_k(x))(\sqrt{x} - P_k(x)) \leq P_k(x) + (\sqrt{x} - P_k(x)) = \sqrt{x}.$$

Hence  $P_{k+1}(x) \leq \sqrt{x}$ . We have therefore shown inductively that  $0 \leq P_k(x) \leq \sqrt{x}$  for every  $k \geq 0$ . As seen above, this also implies that  $P_{k+1} \geq P_k(x)$ .

It follows that for any fixed  $x \in [0, 1]$ ,  $\{P_k(x)\}$  is monotone increasing and bounded above by 1, and hence converges to a limit  $P(x) \leq 1$ . But then  $P(x)$  satisfies

$$P(x) = P(x) + \frac{x - P(x)^2}{2}$$

and hence

$$P(x)^2 = x.$$

Since  $P(x) \geq 0$ , we conclude that  $P(x) = \sqrt{x}$  and  $P_k$  converges pointwise to the square root function. Since the convergence is monotone and the limit function is continuous, Dini's theorem implies that the convergence is actually uniform.

Now let  $\epsilon > 0$ . Pick  $k$  so that  $|P_k(x) - \sqrt{x}| < \epsilon$  for all  $x \in [0, 1]$ . Define  $q(y) = P_k(y^2)$  for  $y \in [-1, 1]$ , so  $q$  is a polynomial. Then for any  $y \in [-1, 1]$ ,

$$|q(y) - \text{abs } y| = |P_k(y^2) - \sqrt{y^2}| < \epsilon$$

since  $y^2 \in [0, 1]$ . Since  $\epsilon > 0$  is arbitrary, we conclude that  $\text{abs} \in P[0, 1]$ .