

Name: _____

Student Id: _____

Calculator Model: _____

Rules:

You have 70 minutes to complete the exam.

Partial credit will be awarded, but you must show your work.

A scientific or graphing calculator is allowed.

A one page sheet of paper (8 1/2 in. x 11 in.) with handwritten notes on one side is allowed.

No other aids are permitted.

Place a box around your **FINAL ANSWER** to each question where appropriate.

If you need extra space, you can use the back sides of the pages. Please make it obvious when you have done so.

Turn off anything that might go beep during the exam.

Good luck!

Problem	Possible	Score
1	10	
2	10	
3	10	
4	10	
5	10	
6	10	
Extra Credit	5	
Total	60	

1. (10 points)

A quantity of a little more than a mole of gas satisfies

$$PV = 9T$$


where pressure P is measured in kPa, volume V is in liters and temperature T is in Kelvin.

a. At time $t = 0$ the temperature of the gas is 300 Kelvin and the container containing it has a volume of 30 liters. What is the pressure of the gas, including units?

b. At time $t = 0$ we know additionally that the temperature of the gas is rising at a rate of $dT/dt = 10$ K/hour and that the volume of the container is decreasing at a rate $dV/dt = -2$ liters/hour. Determine dP/dt , including all units.

2. (10 points)

A thin metal plate has the shape below:



The mass density of the plate is given by $\rho = 5(10 - r)$ grams per square centimeter, where r is the distance in centimeters from the vertex of the plate.

Compute the mass of the plate. **Hint:** the equation we have for area in polar coordinates was for angles in radians.

3. (10 points)


We wish to optimize the function

$$f(x, y) = xe^y$$

subject to the constraint $g(x, y) = x^2 + y^2 = 1$.

a. The figure below depicts level sets of f . Add the following to the figure:

- The gradient of f at the point $(1, 0)$
- The constraint set (i.e. all the points satisfying $g(x, y) = 1$)
- The gradient of g at the point $(0, 1)$



b. In the figure, label **with a square** the location on the constraint set where f is **minimized**. At that point also add to the figure the gradient of f and the gradient of g . Clearly label which gradient goes with which function.

c. Set up the system of equations to solve (using the method of Lagrange multipliers) to determine the location of the minimizer. **Do not solve the system!!**

4. (10 points)

Determine all critical points of the function

$$f(x, y) = y(e^x - 1)$$

and classify each as a local min, local max or saddle.

