

Instructions. You have 90 minutes. Closed book, closed notes, no calculator. *Show all your work* in order to receive full credit.

1. Show that $\lim_{(x,y) \rightarrow (-1,1)} \frac{xy+1}{2x^2-y^2-1}$ does not exist.

Solution:

- Setting $x = -1$ and letting $y \rightarrow 1$ to approach $(-1, 1)$ along the line $(-1, y)$, we see

$$\lim_{y \rightarrow 1} \frac{1-y}{1-y^2} = \frac{1}{2}.$$

- Setting $y = 1$ and letting $x \rightarrow -1$ to approach $(-1, 1)$ along the line $(x, 1)$, we see

$$\lim_{x \rightarrow -1} \frac{x+1}{2x^2-2} = -\frac{1}{4}.$$

Since these limits are different, the original multivariable limit does not exist.

2. Use Lagrange multipliers to find the point(s) on the curve $x^2 - 2y^2 = 1$ closest from the point $P(0, 2)$.

Solution: We want to minimize the distance from a point on the hyperbolic curve to $P(0, 2)$. For simplicity, let $f(x, y)$ be the square of that distance:

$$f(x, y) = (x - 0)^2 + (y - 2)^2 = x^2 + (y - 2)^2.$$

Then our constraint is $g(x, y) = x^2 - 2y^2 = 1$ and we need also to satisfy:

$$\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2x, 2(y-2) \rangle = \lambda \langle 2x, -4y \rangle \quad \Rightarrow \quad \begin{cases} 2x = 2\lambda x \\ 2(y-2) = -4\lambda y \end{cases}$$

The first equation has two solutions:

- either $x = 0$, then from the constraint, $0 - 2y^2 = 1$ which has no real solution for y ;
- or $\lambda = 1$, then from the second equation:

$$2y - 4 = -4y \quad \Rightarrow \quad y = \frac{2}{3}$$

and so plugging into the constraint $x^2 = 1 + 2\left(\frac{4}{9}\right) = \frac{17}{9}$ so we have the points $\left(\pm \frac{\sqrt{17}}{3}, \frac{2}{3}\right)$.

Both have the same $f(x, y)$ value so they are both points we're looking for: $\left(\pm \frac{\sqrt{17}}{3}, \frac{2}{3}\right)$.

3. Find an equation of the tangent plane to the following surface at the point $(x_0, y_0, z_0) = (2, 1, -1)$:

$$x \ln y - 3yz^2 + 1 = xz.$$

Solution: Let $F(x, y, z) = x \ln y - 3yz^2 - xz = -1$. Then,

$$\nabla F(2, 1, -1) = \left\langle \ln y - z, \frac{x}{y} - 3z^2, -6yz - x \right\rangle \Big|_{(2,1,-1)} = \langle 0 + 1, 2 - 3, 6 - 2 \rangle = \langle 1, -1, 4 \rangle$$


and so the equation of the tangent plane is:

$$(x - 2) - (y - 1) + 4(z + 1) = 0 \quad \Rightarrow \quad \boxed{x - y + 4z + 3 = 0}.$$

4. For each of the iterated integrals below, sketch the region of integration then convert as indicated. DO NOT evaluate.

(a) Rewrite $\int_{-2}^0 \int_0^{x^2} 3xy \, dy \, dx$ in the order $dx \, dy$.


Solution:



$$\int_0^4 \int_{-2}^{-\sqrt{y}} 3xy \, dx \, dy$$


(b) Rewrite $\int_{-\pi/2}^{\pi/4} \int_0^1 r^2 \, dr \, d\theta$ in rectangular coordinates.

Solution: From the picture below, we need to split the integral. The order $dx \, dy$ is a bit easier as the split is at $y = 0$ but we still need to solve for y when $\theta = \frac{\pi}{4}$ and $r = 1$, i.e. $y = 1 \sin \frac{\pi}{4} = \frac{\sqrt{2}}{2}$. Inner bounds are from the y -axis $x = 0$, the circle $x^2 + y^2 = 1$, and $y = x$:



$$\int_{-1}^0 \int_0^{\sqrt{1-y^2}} \sqrt{x^2 + y^2} \, dx \, dy + \int_0^{\frac{\sqrt{2}}{2}} \int_y^{\sqrt{1-y^2}} \sqrt{x^2 + y^2} \, dx \, dy$$

5. Compute the mass m of the planar lamina with density $\rho(x, y) = y^2$ shown below.



Solution:

$$\begin{aligned} m &= \iint_R y^2 \, dA = \int_0^{\pi/2} \int_0^2 r^2 \sin^2 \theta \, r \, dr \, d\theta = \int_0^{\pi/2} \left[\frac{r^4}{4} \sin^2 \theta \right]_0^2 \, d\theta = \int_0^{\pi/2} 4 \sin^2 \theta \, d\theta \\ &= \int_0^{\pi/2} 2(1 - \cos(2\theta)) \, d\theta = \left[2\theta - \sin(2\theta) \right]_0^{\pi/2} = \boxed{\pi}. \end{aligned}$$

6. Consider the function:

$$f(x, y) = x^3 - 12xy + 8y^3.$$

(a) Find and classify all critical points of $f(x, y)$.

Solution:

- Find the critical points from solving $\nabla f = \vec{0}$:

$$\nabla f = \vec{0} \quad \Rightarrow \quad \langle 3x^2 - 12y, -12x + 24y^2 \rangle = \langle 0, 0 \rangle \quad \Rightarrow \quad \begin{cases} x^2 = 4y \\ x = 2y^2 \end{cases} \quad \Rightarrow \quad \begin{cases} 4y^4 = 4y \\ x = 2y^2 \end{cases}$$

The first equation simplifies to $y(y^3 - 1) = 0$ so either $y = 0$ or $y = 1$. Substituting back into the second equation gives us the two critical points $(0, 0)$ and $(2, 1)$.

- Apply the Second Partials Test to classify them:

$$f_{xx} = 6x, f_{yy} = 48y, f_{xy} = -12 \Rightarrow d = f_{xx}f_{yy} - f_{xy}^2 = 288xy - 144 = 144(2xy - 1)$$

$d(0, 0) = -144 < 0$ so we have a saddle point at $(0, 0, 0)$;

$d(2, 1) = 144(3) > 0$ and $f_{xx}(2, 1) = 12 > 0$ so f has a local minimum at $(2, 1)$.

- (b) Find the absolute minimum and maximum values of $f(x, y)$ in the rectangular region R defined by $0 \leq x \leq \frac{1}{2}$ and $0 \leq y \leq 1$.

Solution: The absolute min/max can happen only at either the critical points within R or on the boundary of R :

- out of the critical points, only $(0, 0)$ is part of R ;
- we will need to check the vertices $(0, 0)$, $(0, 1)$, $(1/2, 0)$, and $(1/2, 1)$;
- along $x = 0$ for $0 \leq y \leq 1$:

$$g(y) = f(0, y) = 8y^3 \Rightarrow g'(y) = 24y^2$$

and $g'(y) = 0$ for $y = 0$ and we find again $(0, 0)$;

- along $x = 1/2$ for $0 \leq y \leq 1$:

$$g(y) = f(1/2, y) = \frac{1}{8} - 6y + 8y^3 \Rightarrow g'(y) = -6 + 24y^2$$

and $g'(y) = 0$ for $y = \pm \frac{1}{2}$; only $(1/2, 1/2)$ is in R ;

- along $y = 0$ for $0 \leq x \leq \frac{1}{2}$:

$$g(x) = f(x, 0) = x^3 \Rightarrow g'(x) = 3x^2$$

and $g'(x) = 0$ for $x = 0$ and we find again $(0, 0)$;

- along $y = 1$ for $0 \leq x \leq \frac{1}{2}$:

$$g(x) = f(x, 1) = x^3 - 12x + 8 \Rightarrow g'(x) = 3x^2 - 12$$

and $g'(x) = 0$ for $x = \pm 2$; neither points are in R .


We now plug in all values of those points into f to find the absolute min/max:

x	y	$f(x, y)$	
0	0	0	
0	1	8	absolute max
$\frac{1}{2}$	0	$\frac{1}{8}$	
$\frac{1}{2}$	1	$\frac{17}{8}$	
$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{15}{8}$	absolute min

7. Evaluate the following.

(a) the volume below the plane $6x + 3y + 2z = 6$ in the first octant:

Solution:




Rewrite $2z = 6 - 6x - 3y$ so $z = 3 - 3x - \frac{3}{2}y$ and the base is bounded (from setting $z = 0$) by the line: $6x + 3y = 6$, i.e. $2x + y = 2$ for $x, y \geq 0$. So we can write $0 \leq y \leq 2 - 2x$ and in x solve for the upper bound by setting $y = 0$ in the line. Then the volume is:

$$\begin{aligned} V &= \int_0^1 \int_0^{2-2x} 3 - 3x - \frac{3}{2}y \, dy \, dx = \int_0^1 \left[(3 - 3x)y - \frac{3}{4}y^2 \right]_{y=0}^{y=2-2x} \, dx \\ &= \int_0^1 3(1-x)(2-2x) - \frac{3}{4}(2-2x)^2 - 0 \, dx = \int_0^1 6(1-x^2) - 3(1-x)^2 \, dx \\ &= \int_0^1 3(1-x)^2 \, dx = \left[-(1-x)^3 \right]_0^1 = 0 + 1 = \boxed{1}. \end{aligned}$$

(b) the surface area of the cone $z = \sqrt{x^2 + y^2}$ above the region R bounded by the graphs of $y = -x$, $x = 2y - y^2$, $y = 0$ and $y = 1$ as sketched below:


Solution: The gradient is $\nabla z = \langle z_x, z_y \rangle = \left\langle \frac{x}{\sqrt{x^2 + y^2}}, \frac{y}{\sqrt{x^2 + y^2}} \right\rangle$ so noting that R is horizontally simple, we have that the surface area of the cone above R is:



$$\begin{aligned} SA &= \iint_R \sqrt{1 + z_x^2 + z_y^2} \, dA = \int_0^1 \int_{x=-y}^{x=2y-y^2} \sqrt{1 + \frac{x^2}{x^2 + y^2} + \frac{y^2}{x^2 + y^2}} \, dx \, dy = \sqrt{2} \int_0^1 \int_{-y}^{2y-y^2} \, dx \, dy \\ &= \sqrt{2} \int_0^1 [x]_{-y}^{2y-y^2} \, dy = \sqrt{2} \int_0^1 2y - y^2 + y \, dy = \sqrt{2} \int_0^1 3y - y^2 \, dy \\ &= \sqrt{2} \left[\frac{3y^2}{2} - \frac{y^3}{3} \right]_0^1 = \sqrt{2} \left(\frac{3}{2} - \frac{1}{3} - 0 \right) = \boxed{\frac{7\sqrt{2}}{6}}. \end{aligned}$$

- (c) the volume of the solid bounded by the paraboloid $z = x^2 + y^2$ and the inverted cone $z = 6 - \sqrt{x^2 + y^2}$ using polar coordinates.

Solution: The cone is above the paraboloid and for the base, we have a disk where the radius can be found using the intersection of the surfaces, i.e. set $x^2 + y^2 = 6 - \sqrt{x^2 + y^2}$ or in polar $r^2 = 6 - r$ for $r = \sqrt{x^2 + y^2} \geq 0$. So $r^2 + r - 6 = 0$ which has for solutions $r = -3, 2$ and we keep $r = 2$. And so the volume is:



$$\begin{aligned}
 V &= \int_0^{2\pi} \int_0^2 (6 - r - r^2) r \, dr \, d\theta = \int_0^{2\pi} \int_0^2 6r - r^2 - r^3 \, dr \, d\theta \\
 &= \int_0^{2\pi} \left[3r^2 - \frac{r^3}{3} - \frac{r^4}{4} \right]_0^2 \, d\theta = \int_0^{2\pi} 12 - \frac{8}{3} - 4 - 0 \, d\theta = \frac{16}{3} [\theta]_0^{2\pi} = \boxed{\frac{32\pi}{3}}.
 \end{aligned}$$

8. The bee population in a boxed beehive is given at each point (x, y, z) by

$$f(x, y, z) = x^2 + y^2 + xyz.$$

- (a) At the point $(3, 1, 2)$, what is the unit direction of greatest decrease in population?

Solution:

$\nabla f(3, 1, 2) = \langle 2x + yz, 2y + xz, xy \rangle|_{(3,1,2)} = \langle 8, 8, 3 \rangle$, so the unit direction of greatest decrease is

$$\boxed{-\frac{\nabla f(3, 1, 2)}{\|\nabla f(3, 1, 2)\|} = \left\langle -\frac{8}{\sqrt{137}}, -\frac{8}{\sqrt{137}}, \frac{3}{\sqrt{137}} \right\rangle.}$$

- (b) Find the directional derivative of f at $(3, 1, 2)$ in the direction of $\mathbf{v} = \langle 1, 2, 2 \rangle$?

Solution:

The direction we consider is $\mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|}$, so $\mathbf{u} = \langle 1/3, 2/3, 2/3 \rangle$. Then

$$D_{\mathbf{u}}f(3, 1, 2) = \nabla f(3, 1, 2) \cdot \mathbf{u} = \langle 8, 8, 3 \rangle \cdot \langle 1/3, 2/3, 2/3 \rangle = \frac{8}{3} + \frac{16}{3} + \frac{6}{3} = \boxed{10}.$$

- (c) Use the chain rule (no direct substitution) to find $\frac{df}{dt}$ in terms of t if $x(t) = 4 - t^2$, $y(t) = 3t - 2$ and $z(t) = 3t^3 - 1$.

Solution:

$$\begin{aligned}
 \frac{df}{dt} &= \nabla f \cdot \left\langle \frac{dx}{dt}, \frac{dy}{dt}, \frac{dz}{dt} \right\rangle = \langle 2x + yz, 2y + xz, xy \rangle \cdot \langle -2t, 3, 9t^2 \rangle \\
 &= (2x + yz)(-2t) + (2y + xz)(3) + (xy)(9t^2) \\
 &= -2t(2(4 - t^2) + (3t - 2)(3t^3 - 1)) + 3(2(3t - 2) + (4 - t^2)(3t^3 - 1)) + 9t^2(4 - t^2)(3t - 2) \\
 &= -2t(9t^4 - 6t^3 - 2t^2 - 3t + 10) + 3(-3t^5 + 12t^3 + t^2 + 6t - 8) + 9t^2(-3t^3 + 2t^2 + 12t - 8) \\
 &= -18t^5 + 12t^4 + 4t^3 + 6t^2 - 20t - 9t^5 + 36t^3 + 3t^2 + 18t - 24 - 27t^5 + 18t^4 + 108t^3 - 72t^2 \\
 &= \boxed{-54t^5 + 30t^4 + 148t^3 - 63t^2 - 2t - 24}
 \end{aligned}$$