

## Derivatives + Rates of Change.

Suppose a ball is tossed in the air




$y(t)$  is height of ball, in m.  $t$  in s.

$$y(0) = 0, \text{ say}$$

$$y(t) = 15t - 5t^2$$

$$y(0) = 0 \quad y(3) = 15 \cdot 3 - 5 \cdot 3 \cdot 3 = 0$$



$$y(1) = 10$$

$$y(2) = 10$$


30-20 ✓

Average rates of change of height:

$$[0, 1] \quad \frac{y(1) - y(0)}{1 - 0} = 10 \text{ m/s}$$

$$[1, 2] \quad \frac{y(2) - y(1)}{2 - 1} = \frac{10 - 10}{1} = 0 \text{ m/s}$$

$$[2, 3] \quad \frac{y(3) - y(2)}{3 - 2} = \frac{0 - 10}{1} = -10 \text{ m/s}$$



The slopes of the three secant lines equal the average rates of change over the three time intervals.

What is the instantaneous rate of change in height (velocity) at  $t=1$ ?

We can estimate: 10 or 0, or better yet  $\frac{10+0}{2}$ .

To formalize:

$$\frac{h(1+h) - h(1)}{h}$$



interpret as either

a) avg rate of change over  $[1, 1+h]$

b) slope of secant line.

As we take the limit as  $h \rightarrow 0$   
we pick up the instantaneous rate of change of height  
(AKA velocity)

or the slope of the tangent line

15k-5k<sup>2</sup>


Let's do it:

$$\begin{aligned}y(1+h) &= 15(1+h) - 5(1+h)^2 \\ &= 15 + 15h - 5(1+2h+h^2) \\ &= 10 + 15h - 10h - 5h^2 \\ &= 10 + 5h - 5h^2\end{aligned}$$

$$y(1) = 15 \cdot 1 - 5 \cdot 1^2 = 10$$

$$\frac{y(1+h) - y(1)}{h} = \frac{5h - 5h^2}{h} \quad \text{but you can't set } h=0.$$

$$\text{But } \lim_{h \rightarrow 0} \frac{5h - 5h^2}{h} = \lim_{h \rightarrow 0} 5 - 5h = \boxed{5}$$



The velocity at  $t=1$  is 5 m/s.

The slope of the tangent line is 5.

What's the equation of the tangent line?

Form  $y = mx + b$     blech.

$$y - y_i = m(x - x_i)$$

point slope!  
If you know  $m$ ,  $y_i$ ,  $x_i$   
good to go.

$$x_i = 1, \quad y_i = 10$$

$$y - 10 = 5(x - 1)$$

$$y = 5x + 5$$

slope 5? ✓  
(1,10) ? ✓

This limit

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

is known as the

derivative of  $f$  at  $a$

and has the notation  $f'(a)$ .

Alternatively  $b = a+h$  ( $h = b-a$ )

as  $h \rightarrow 0$ ,  $b \rightarrow a$ .

$$\lim_{b \rightarrow a} \frac{f(b) - f(a)}{b - a}$$

You need to know both expressions.

What does the derivative tell you?

Think of population at time  $t$

$$\frac{P(b) - P(a)}{b - a} \quad \frac{\Delta P}{\Delta t}$$

If  $b$  is close to  $a$ ,  $\frac{\Delta P}{\Delta t} \approx P'(a)$

$$\Delta P \approx P'(a) \Delta t$$

If you change time by a little bit  $\Delta t$ ,  
the population will approximately  
change by  $P'(a) \Delta t$

E.g. if  $P'(3) = 122 \frac{\text{units}}{\text{year}}$  ( $t$  in years)

then over 6 months, pop will change by

$$\Delta P = P'(3) \Delta t = 122 \frac{\text{units}}{\text{year}} \cdot \frac{1}{2} \text{ year} = 61 \text{ units per year}$$

Now you:

The radius of a tree is given by

$$r(t) = 10\sqrt{t} \quad r \text{ in cm, } t \text{ in years.}$$

a) Find the average rate of change of growth from  $t=1$  to  $t=2$ .

b) Find the instant rate of change of growth at  $t=1$ .

c) I promise you

$$r(4) = 20$$

$$r'(4) = 5/2$$

$$10 \int \frac{1}{\sqrt{t}}$$

use these facts alone to approximate  $r$  at 4 years and 1 month.

$$20 + \frac{5}{2} \cdot \frac{1}{12}$$

then compare your est.