

Other trig derivatives:

$$\frac{d}{dx} \tan(x) = \frac{d}{dx} \frac{\sin(x)}{\cos(x)} = \sec^2(x) \quad (\text{You did these!})$$

$$\frac{d}{dx} \cot(x) = \frac{d}{dx} \frac{\cos(x)}{\sin(x)} = \frac{-\sin^2(x) - \cos^2(x)}{\sin^2(x)} = -\csc^2(x)$$

$$\frac{d}{dx} \sec(x) = \frac{d}{dx} \frac{1}{\cos(x)} = -\frac{-\sin(x)}{\cos^2(x)} = \frac{\sin(x)}{\cos(x)} \frac{1}{\cos(x)} = \sec(x) \tan(x)$$

$$\frac{d}{dx} \csc(x) = \frac{d}{dx} \frac{1}{\sin(x)} = \frac{-\cos(x)}{\sin^2(x)} = -\cot(x) \csc(x)$$


2. If you increase the radius of a snowball from 2 inches to 2.02 inches, estimate the change in volume of the snowball.

$$\Delta r = 2.02 - 2 = 0.02.$$

$$\Delta V \approx V'(2) \cdot \Delta r = 50.265 \cdot (0.02) = 1.0053 \text{ cubic inches}$$

6. A 12 foot ladder rests against a wall. Let θ be the angle between the ladder and the wall and let x be the distance from the base of the ladder and the wall.

- a. Compute x as a function of θ .



$$\sin \theta = \frac{x}{12}$$

$$x = 12 \sin \theta$$


- b. How fast does x change with respect to θ when $\theta = \pi/6$?

$$x(\theta) = 12 \sin \theta$$

$$x'(\theta) = 12 \cos \theta$$

$$x'\left(\frac{\pi}{6}\right) = 12 \cos \frac{\pi}{6}$$

$$= 12 \cdot \frac{1}{2} = 6.$$



Chain Rule

We've seen: $\frac{d}{dx} e^{2x} = 2e^{2x}$

$$\frac{d}{dx} e^{-x} = -e^{-x}.$$

There is a pattern here. To see more, consider

$$\begin{aligned}\frac{d}{dx} \sin(5x) &= \lim_{h \rightarrow 0} \frac{\sin(5(x+h)) - \sin(5x)}{h} \\ &= 5 \lim_{h \rightarrow 0} \frac{\sin(5x + 5h) - \sin(5x)}{5h} \\ &= 5 \lim_{w \rightarrow 0} \frac{\sin(5x + w) - \sin(5x)}{w} \\ &= 5 \sin'(5x) \\ &= 5 \cos(5x)\end{aligned}$$

In general $\frac{d}{dx} \sin(ax) = a \cos(ax)$

$$\frac{d}{dx} e^{ax} = a e^{ax}$$

These are special cases of the following rule, known as the Chain Rule

$$\frac{d}{dx} f(g(x)) = f'(g(x)) g'(x)$$

$$\begin{aligned} \frac{d}{dx} \sin(5x) &= \sin'(5x) \cdot \frac{d}{dx}(5x) \\ &= 5 \cos(5x) \end{aligned}$$

$$\begin{aligned} \text{e.g. } \frac{d}{dx} \cos(x^2) &= -\sin(x^2) \frac{d}{dx} x^2 \\ &= -2x \sin(x^2) \end{aligned}$$

Now your turn, Worksheet-3-4.