

5. For each of the following functions, find a particular antiderivative.

Function	Antiderivative	Function	Antiderivative
x		$\sin(x)$	
x^2		$\cos(x)$	
x^3		e^x	
$x^k (k \neq -1)$		$1/(1+x^2)$	
x^{-1} for $x > 0$		$\sec^2(x)$	
x^{-1} for $x < 0$		$\sec(x) \tan(x)$	
x^{-1} for all x		1	

6. Compute three different antiderivatives of $f(x) = x^{20} + 4x^{10} + 8$

7. Compute an antiderivative of $f(t) = \frac{5 \sec t \tan t}{3} - 4 \sin t - \frac{1}{t} + e^2$

8. Compute an antiderivative of $f(x) = \cos(3x)$.

9. Compute the antiderivative of $f(t) = t^2$ that equals 5 when $t = 2$.
10. A particle moves in a straight line and has acceleration given by $a(t) = 5 \cos t - 2 \sin t$. Its initial velocity is $v(0) = -6$ m/s and its initial position is $s(0) = 2$ m. Find its position function $s(t)$.

11. A stone is dropped from a cliff and hits the ground three seconds later. How high is the cliff? (Acceleration due to gravity is 9.8 m/s^2 .)

12. What constant acceleration is needed to take a car from 10 mph to 60 mph in 5 seconds?