

$$= \lim_{h \rightarrow 0} \frac{\sin(x) \cos(h) + \sin(h) \cos(x) - \sin(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\sin(x) [\cos(h) - 1] + \sin(h) \cos(x)}{h}$$

$$= \sin(x) \lim_{h \rightarrow 0} \frac{\cos(h) - 1}{h} + \cos(x) \lim_{h \rightarrow 0} \frac{\sin(h)}{h}$$

$$= \sin(x) \cdot 0 + \cos(x) \cdot \underline{1}$$

$$= \cos(x) \left[ \frac{d}{dt} \cos(x) \right]$$

# Section 3.5 (Implicit Differentiation)

$$\frac{d}{dx} \ln(x)$$

$$\frac{d}{dx} \arctan(x)$$

$$f(x) = x^{1/3}$$

$$\rightarrow \left[ \frac{1}{3} x^{-2/3} \right]$$



$$\frac{df}{dx}$$

$$y = x^{1/3}$$

$$f'(x)$$

$$y = x^{1/3}$$

$$y^3 = x$$



$$\frac{dy}{dx}$$

$$y = x^{1/3}$$

$$y^3 = x$$

$$\rightarrow (y(x))^3 = x$$

↓

$$\frac{d}{dx} (y(x))^3 = \frac{d}{dx} x = 1$$

||

$$3 (y(x))^2 \cdot \frac{d}{dx} y(x)$$

$$3 (y(x))^2 \frac{dy}{dx} = 1$$

$$\frac{d}{dt} (\sin(t))^3 = 3 \sin^2(t) \cdot \cos(t)$$

$$\frac{dy}{dx} = \frac{1}{3(y(x))^2}$$

$$y(x) = x^{1/3}$$

$$\frac{dy}{dx}$$

$$\frac{dy}{dx} = \frac{1}{3(x^{1/3})^2} = \frac{1}{3}$$

$$\frac{1}{x^{2/3}} = \frac{1}{3} x^{-2/3}$$

$$\frac{dy}{dx} = \frac{1}{3} x^{-2/3}$$

Find  $\frac{dy}{dx}$  if  $y = x^{1/3}$ .

Start with  $y^3 = x$

$$\frac{d}{dx} y^3 = \frac{d}{dx} x$$

$$3y^2 \cdot \frac{dy}{dx} = 1$$

$$y = x^{2/3}$$

$$y^2 = x^{4/3}$$


$$\frac{dy}{dx} = \frac{1}{3y^2}$$

$$\frac{dy}{dx} = \frac{1}{3} \frac{1}{x^{2/3}} = \frac{1}{3} x^{-2/3}$$

Silly geometric fun:

$$y = \sqrt{17 - x^2}$$

Consider the curve  $x^2 + y^2 = 17$



$$(x, y) = (1, 4)$$

is on the curve.

We want to compute the equation of the tangent line to the curve at  $(1, 4)$

We need the slope of the tangent line.

We want  $\frac{dy}{dx}$  at  $x=1$  ( $y=4$ )

$$x^2 + y^2 = 1$$

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx} 1$$

$$\frac{d}{dx} x^2 + \frac{d}{dx} y^2 = 0$$

$$\hookrightarrow (y(x))^2$$

$$\frac{d}{dx} (y(x))^2 = 2y(x) \frac{d}{dx} y(x)$$

$$2x + 2y \cdot \frac{dy}{dx} = 0$$


$$2y \frac{dy}{dx} = -2x$$

$$\frac{dy}{dx} = \frac{-2x}{2y} = -\frac{x}{y}$$

$$x^2 + y^2 = 17$$

$$\frac{dy}{dx} = -\frac{x}{y}$$

$$\frac{dy}{dx} = -\frac{1}{4} \text{ at } (1, 4)$$



$$(-4, 1) = (x, y)$$

$$x^2 + y^2 = 17$$

$$\frac{dy}{dx} = -\frac{-4}{1} = +4$$

---

$$x^3 = y^4 + x^2 \sin(y) + 1$$

← curve

$$1^3 = 0^4 + 1^2 \sin(0) + 1$$

point on the curve

$$(x, y) = \underline{(1, 0)}$$

What is  $\frac{dy}{dx}$  at this point?



$$(x, y) = (1, 0)$$

$$x^3 = y^4 + x^2 \sin(y) + 1$$

$$\frac{d}{dx} x^3 = \frac{d}{dx} \left[ y^4 + x^2 \sin(y) + 1 \right]$$

$$3x^2 = \frac{d}{dx} y^4 + \frac{d}{dx} (x^2 \sin(y)) + \frac{d}{dx} 1$$

$$= 4y^3 \frac{dy}{dx} + \frac{d}{dx} (x^2) \cdot \sin(y) + x^2 \frac{d}{dx} \sin(y)$$

$$= 4y^3 \frac{dy}{dx} + 2x \sin(y) + x^2 \cos(y) \frac{dy}{dx}$$

$$3x^2 = 4y^3 \frac{dy}{dx} + 2x \sin(y) + x^2 \cos(y) \frac{dy}{dx}$$

$$3x^2 - 2x \sin(y) = \left[ 4y^3 + x^2 \cos(y) \right] \frac{dy}{dx}$$

$$\frac{dy}{dx} = \frac{3x^2 - 2x \sin(y)}{4y^3 + x^2 \cos(y)}$$

$$x^3 = y^4 + x^2 \sin(y) + 1 \quad (1, 0)$$


$$\text{At } (x, y) = (1, 0)$$

$$\frac{dy}{dx} = \frac{3 \cdot 1^2 - 2 \cdot 1 \sin(0)}{4 \cdot 0^3 + 1^2 \cos(0)} = \frac{3 - 2 \cdot 0}{0 + 1}$$

$$= \frac{3}{1}$$


$$= 3$$

# Inverse Trig Functions



$$\frac{\sin(\theta)}{\cos(\theta)}$$

$$\tan(\theta) = 17$$



$$\tan(\arctan(z)) = z \quad \text{always}$$

$$\arctan(\tan(\theta)) = \theta \quad \text{only if}$$

$$-\frac{\pi}{2} < \theta < \frac{\pi}{2}$$

Notation

$$\tan^{-1}(x)$$



~~$$\tan(x)$$~~



Ush.