

1. Recall the two step Adams-Bashforth method:

$$u_{i+1} = u_i + \frac{h}{2}(3f_i - f_{i-1}).$$

- a) Write down the stability polynomial $p(\rho)$ for this method applied to the problem $u' = \lambda u$.
- b) The equation $p(\rho) = 0$ will involve the expression λh . Solve for λh to write

$$\lambda h = f(\rho)$$

for some function f .

- c) Make a numerical sketch of the boundary of the region of absolute stability as follows. The characteristic polynomial has the form $p(\rho) = \sigma(\rho) - z\eta(\rho)$ for certain polynomials σ and η . The boundary of the region of absolute stability is given by those values of z such that there is a root of $p(\rho)$ with $|\rho| = 1$. Turn this idea on its head: the condition $p(\rho) = 0$ is the same as

$$z = \frac{\sigma(\rho)}{\eta(\rho)}.$$

So if you plug in a ρ on the right-hand side with $|\rho| = 1$, you get a z such that the characteristic polynomial has a root with size 1. You can generate a curve of all such values of z by letting ρ vary around the unit circle of the complex plane.

2. Recall that a linear multistep method (LMM) has the form

$$\alpha_k u_{k+n} + \cdots + \alpha_1 u_{1+n} + \alpha_0 u_n = h(\beta_k f_{k+n} + \cdots + \beta_1 f_{1+n} + \beta_0 f_n)$$

- a) Show that the method is consistent if and only if

$$\begin{aligned} \alpha_k + \cdots + \alpha_0 &= 0 \\ k\alpha_k + \cdots + 1\alpha_1 + 0\alpha_0 &= \beta_k + \cdots + \beta_0 \end{aligned} \tag{1}$$

- b) Use the previous result to show that every consistent one-step LMM is zero stable.

3. Implement Euler's method and the Runge-Kutta RK4 method described in Table 1.3 for **vector** valued ODEs (i.e, systems). Test your work against the IVP $\mathbf{x}' = A\mathbf{x}$ where

$$A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

with initial condition $\mathbf{x}_0 = (1, 0)$. Part of the exercise is to compute the true solution of this system. Hint: convert into a second-order scalar ODE. **Verify that your code has the theoretical order of convergence.**

4. You are going to solve the 2-body problem of gravitation for the Earth-moon system. Give two bodies with masses m_1 and m_2 at positions x_1 and x_2 in cartesian coordinates, the force of body 2 on body 1 is

$$F_{21} = Gm_1m_2 \frac{x_2 - x_1}{|x_2 - x_1|^3}$$

where G is the gravitational constant. The force of body 1 on body 2 is the same, with the numbers 1 and 2 interchanged.

Newton's Laws then read

$$\begin{aligned} m_1 \mathbf{x}_1'' &= F_{21} \\ m_2 \mathbf{x}_2'' &= F_{12}. \end{aligned} \tag{2}$$

We have the following physical constants:

$$m_{\text{Earth}} = 5.972 \times 10^{24} \text{ kg}$$

$$m_{\text{moon}} = 7.342 \times 10^{22} \text{ kg}$$

$$G = 6.67408 \times 10^{-11} \frac{\text{m}^3}{\text{kg s}^2}$$

- a) Convert system (2) into a first order system by introducing the variables $\mathbf{v}_1 = \mathbf{x}_1'$ and $\mathbf{v}_2 = \mathbf{x}_2'$.
- b) Writing $\mathbf{x}_i = [x_i, y_i]$ and $\mathbf{v}_i = [v_i, w_i]$, we will keep track of all of the scalar system variables in vectors

$$[x_1, y_1, x_2, y_2, v_1, w_1, v_2, w_2]^T.$$

Using this convention, write a right-hand side function $\mathbf{z} = \text{earthmoon}(\mathbf{t}, \mathbf{u})$ for the system you wrote down in part 1.

- c) Suppose at time $t = 0$ the Earth is stationary and located at the origin, the moon has position $x = 3.565 \times 10^8$ and $y = 0$ meters with velocity $1.09 \times 10^3 \text{ m/s}$ in the positive y direction.

Use these initial conditions, and each of the solvers from problem 2, to generate an approximate solutions over a 40 day time interval (convert to seconds!) using daily ($M=40$) and then hourly ($M=40 \times 24$) time steps. That is, you are generating four different approximate solutions. Make basic plots of the computed trajectories. Describe in a few words what you see, and how these results relate to the local truncation errors of the schemes.

- d) An important property of an isolated physical system is that its energy is conserved. For the 2-body problem, the energy is

$$E(t) = \frac{m_1}{2} |\mathbf{x}_1'|^2 + \frac{m_2}{2} |\mathbf{x}_2'|^2 + U(\mathbf{x}_1, \mathbf{x}_2)$$

where U is the potential energy

$$U(\mathbf{x}_1, \mathbf{x}_2) = -\frac{Gm_1m_2}{|\mathbf{x}_1 - \mathbf{x}_2|}.$$

If you have not seen this before, you should compute the derivative $E'(t)$ and show, using the differential equation, that it is zero (and hence $E(t)$ is constant).

Start by computing the value $E(0)$ exactly from the initial conditions. Then compute, for each of your four approximate solutions, an array of scalar energy values $E(t)$ computed at the $M + 1$ sample times. Plot the four energy curves from the four runs in one figure. Explain and comment. Describe an “energy error norm” which is small if the solution is of high accuracy, and report the values for the four runs.