

This lab concerns fitting a polynomial to a number of data points. The most basic version of this operation is fitting a line to two points, a task that is old hat to you. If you have a third point, you can't fit a line to this data, but presumably a quadratic would work. A polynomial that passes through given data points is called a **polynomial interpolant** of the data.

The lab has a companion Jupyter notebook. Follow the instructions below and update the corresponding sections of the notebook as needed. For some problems, you will be asked to attach a hand computation to your final output. To do this, you will make a PDF of your Jupyter notebook and then attach to the end of it scanned PDF pages of your handwritten work.

**Exercise 1:**

To begin, let's think about that line fitting problem again. Suppose we have two points  $p_1 = (x_1, y_1)$  and  $p_2 = (x_2, y_2)$  and we would like to find the equation of a line  $y = mx + b$  going through those points. Substitute these points into the equation of the line to obtain two equations for the unknowns  $m$  and  $b$ .

**Exercise 2:**

Suppose  $p_1 = (0.2, 0.7)$  and  $p_2 = (0.8, -0.4)$ . What are the equations for  $m$  and  $b$ ?

**Exercise 3:**

Write these equations in matrix form:

$$A \begin{bmatrix} b \\ m \end{bmatrix} = v.$$

Explicitly write down what  $A$  and  $v$  are.

**Exercise 4:**

Julia has a built-in facility for solving linear systems of equations expressed in matrix form. Follow the instructions in the notebook to solve the system you wrote down in Exercise 3.

**Exercise 5:**

Verify the  $m$  and  $b$  you just computed work by plotting the line  $y = mx + b$  as well as the two data points that defined the line. The notebook has more information on this task.

**Exercise 6:**

Now find the equation of a parabola  $y = ax^2 + bx + c$  passing through the points  $(-1, 1.5)$ ,  $(3, 32.2)$ ,  $(5, -42.6)$ . You must

- Record, by hand, the system of equations to solve.
- Convert the system into a matrix system.
- Solve the system using Julia (record the command you used and the solution).
- Generate a plot that contains the parabola and the three points. Your  $x$  coordinate on your plot should range from  $x = -2$  to  $x = 6$ .

**Exercise 7:**

If you have 7 data points  $(x_n, y_n)$ , with all of the  $x_n$ 's different, these uniquely determine a polynomial of some order. What is the order? Enter your response in the notebook.

**Exercise 8:**

We're going to want to evaluate polynomials with given coefficients and given points. Write a function `poly_eval` in Julia that receives a vector of polynomial coefficients  $c = (c_0, \dots, c_n)$  and a vector of  $x$ -values,  $x = (x_1, \dots, x_k)$  and returns a vector of polynomial values  $(p(x_1), \dots, p(x_k))$ .

Follow the instructions in the notebook to write this function and test it.

**Exercise 9:**

For a general polynomial  $p(x) = c_0 + c_1x + c_2x^2 + \dots + c_nx^n$ , if you have  $n + 1$  data points  $(x_k, y_k)$  and you want  $p(x_k) = y_k$ , then you obtain  $n + 1$  equations:

$$c_0 + c_1x_k + c_2x_k^2 + \dots + c_nx_k^n = y_k$$

Equivalently, the coefficients  $c = (c_0, c_1, \dots, c_n)$  satisfy

$$Ac = y$$

where  $y = (y_1, \dots, y_{n+1})$  and where

$$A = \begin{bmatrix} 1 & x_1 & \dots & x_1^{n-1} & x_1^n \\ 1 & x_2 & \dots & x_2^{n-1} & x_2^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n+1} & \dots & x_{n+1}^{n-1} & x_{n+1}^n \end{bmatrix}.$$

The matrix  $A$  is called a Vandermonde matrix. Your task: Write a function in Julia that receives a single vector  $x = (x_1, \dots, x_{n+1})$  and returns the associated Vandermonde matrix. Your experience on the Homework 6 Julia problem writing Toeplitz matrices will be helpful! Do that first, if you have not done so already.

**Exercise 10:**

Suppose we have data points  $(1, 4), (2, -1), (4, 7), (5, -3), (8, 1), (9, -10), (11, 3)$ . Use your 'vandermonde' function to obtain the associated Vandermonde matrix  $A$ . Then use Julia to solve  $Ac = y$ . Finally, plot the the polynomial along with the data points that were used to determine it.

**Exercise 11:****Challenge Problem, not due**

Suppose we want to make a polynomial approximation of  $\cos(x)$ . We know the values of  $\cos(x)$  exactly at  $x = 0, \pi/6, \pi/4, \pi/3, \pi/2, 2\pi/3, 3\pi/4, 5\pi/6, \pi$ . We also know that cosine is an even function, so our polynomial should only involve terms of even order. Use Julia to find coefficients  $c_0, \dots, c_8$  such that the polynomial

$$p(x) = c_0 + c_1x^2 + c_2x^4 + \dots + c_8x^{16}$$

satisfies  $p(x) = \cos(x)$  for each value of  $x$  in the list above. Then generate two plots. The first shows  $p(x)$  and the data points it interpolates. The second shows the error  $|p(x) - \cos(x)|$  over the range  $0 \leq x \leq \pi$ . What is the maximum error?