Instructions: (10 points total) Show all work for credit. You may use your book, but no other resource.

- 1. (5 pts.) Consider the solid E which, in cylindrical coordinates, is bounded by the planes $z=0, z=r\sin(\theta)+5$ and the cylinders r=1 and r=5
 - (a) Sketch (as best you can) the solid E.
 - (b) Compute the definite integral $\iiint_E x y \, dV$
 - 1. (5 pts.) Consider the solid E which, in cylindrical coordinates, is bounded by the planes $z=0, z=r\sin(\theta)+5$ and the cylinders r=1 and r=5
 - (a) Sketch (as best you can) the solid E.
 - (b) Compute the definite integral $\iiint_E x y \, dV$

$$= \int_{0}^{2\pi} \int_{0}^{5} (r\cos\theta - r\sin\theta) (r\sin\theta + s) drd\theta$$

$$= \int_{0}^{2\pi} \int_{0}^{5} r^{2} (r\sin\theta + s) (\cos\theta - \sin\theta) drd\theta = \int_{0}^{2\pi} (\cos\theta - \sin\theta) \int_{0}^{5} r^{3} \sin\theta + 5r^{2} dr d\theta$$

$$= \left(\frac{2\pi}{6} \left(\cos \theta - \sin \theta\right) \left[\frac{1}{4} + 4\sin \theta + \frac{5}{3} + \frac{3}{3}\right] + d\theta = \int_{0}^{2\pi} \left(\cos \theta - \sin \theta\right) \left[\frac{625}{4} \sin \theta + \frac{625}{3}\right] - \left(\frac{1}{4} \sin \theta + \frac{5}{3}\right) d\theta$$


$$= \int_{0}^{2\pi} (\cos \theta - \sin \theta) \left(156 \sin t + \frac{620}{3} \right) d\theta = \int_{0}^{2\pi} 156 \cos t \sin t + \frac{620}{3} \cos \theta - 156 \sin^{2} t$$

$$= \int_{0}^{2\pi} (\cos \theta - \sin \theta) \left(156 \sin t + \frac{620}{3} \cos \theta - 156 \sin^{2} t + \frac{620}{3} \cos \theta - \frac{620}{3} \sin t + \frac{620}{3} \cos \theta - \frac{620}{3} \sin t + \frac{620}{3} \cos \theta - \frac{620}{3} \sin t + \frac{620}{3} \cos \theta - \frac{620}{$$

$$= \frac{156 \sin^2 t}{2} + \frac{620}{3} \sin t - \frac{156}{5} \left[\frac{9}{2} - \frac{\sin^2 \theta}{4} \right] + \frac{(20)}{3} \cos t \Big|_{0}^{27}$$

$$= 78 \sin^{2} t \Big|_{0}^{2\pi} - \frac{620}{3} \sin t \Big|_{0}^{2\pi} - \frac{1569}{2} \Big|_{0}^{2\pi} + \frac{1565 \sin 20}{4} \Big|_{0}^{2\pi} + \frac{620}{3} \cos t \Big|_{0}^{2\pi}$$

(a) Without doing any calculus at all, compute the volume of the solid B. (You may look up the volume of a sphere if you do not remember it.)



(b) Now use spherical coordinates and an appropriate triple integral to compute this volume.

(Lots of variants!)
$$\frac{\pi}{2} \pm \theta = 2\pi$$
, $\alpha = \Gamma \pm 4$, $\alpha = \varphi \in \pi/2$

$$= \int_{\pi/2}^{2\pi} \int_{2}^{4} -\rho^{2} \cos \varphi \left(\int_{0}^{\pi/2} dy d\theta \right) = \int_{\pi/2}^{2\pi} \int_{2}^{4} -\rho^{2} \left(o-1 \right) dy d\theta = \int_{\pi/2}^{2\pi} \int_{2}^{4} \rho^{2} d\rho d\theta$$

$$= \int_{\pi_{12}}^{2\pi} \frac{1}{3} \rho^{3} \Big|_{2}^{4} d\theta = \int_{\pi_{12}}^{2\pi} \frac{64}{3} - \frac{8}{3} d\theta = \int_{\pi_{12}}^{2\pi} \frac{56}{3} d\theta = \frac{56}{3} \Big(2\pi - \frac{\pi}{2} \Big)$$

$$=\frac{56}{3}(\frac{3\pi}{2})=\boxed{28\pi}$$